arXiv:0809.2870v1 [math-ph] 17 Sep 2008

Exact solutions for the general fifth order KdV equation by the extended tanh method

Alvaro Salas [*]

Cesar A. Gómez S [†]

José Gonzalo Escobar Lugo [‡]

Abstract

In this paper we show some exact solutions for the general fifth order KdV equation $u_t + \omega\, u_{xxxxx} + \alpha\, uu_{xxx} + \beta u_x u_{xx} + \gamma u^2 u_x = 0$. These solutions are obtained by the extended tanh method.

Key words and phrases: Extended tanh method, nonlinear ordinary differential equation, nonlinear partial differential equation, fifth order evolution equation, fifth order KdV equation, fKdV, traveling wave solution, extended tanh method, partial differential equation (PDE), KdV equation, Caudrey-Dodd-Gibbon equation, Ito equation, Caudrey-Dodd-Gibbon equation, Sawada-Kotera equation, Kaup-Kupershmidt equation, nonlinear evolution equation.

---

[*] Department of Mathematics, Universidad de Caldas, Department of Mathematics, Universidad Nacional de Colombia, Manizales, Colombia. *email* : asalash2002@yahoo.com

[†] Universidad Nacional de Colombia, Bogotá. *email* : cagomezsi@unal.edu.co

[‡] Universidad Cooperativa de Colombia *email* : jogoel@gmail.com

1 Introduction

A large variety of physical, chemical, and biological phenomena is governed by nonlinear evolution equations. The analytical study of nonlinear partial differential equations was of great interest during the last decades. Investigations of traveling wave solutions of nonlinear equations play an important role in the study of nonlinear physical phenomena. The importance of obtaining the exact solutions, if available, of those nonlinear equations facilitates the verification of numerical solvers and aids in the stability analysis of solutions.

The nonlinear generalized KdV equation of fifth order (fKdV equation) reads

$$u_t + \omega\, u_{xxxxx} + \alpha\, u u_{xxx} + \beta\, u_x u_{xx} + \gamma u^2 u_x = 0, \tag{1.1}$$

where α, β, γ and ω are arbitrary real parameters.

This equation describes motions of long waves in shallow water under gravity and in a one-dimensional nonlinear lattice and it is an important mathematical model with wide applications in quantum mechanics and nonlinear optics. Typical examples are widely used in various fields such as solid state physics, plasma physics, fluid physics and quantum field theory. A great deal of research work has been invested during the past decades for the study of the fKdV equation. The main goal of these studies was directed towards its analytical and numerical solutions. Several different approaches, such as Blackund transformation, a bilinear form, and a Lax pair, have been used independently by which soliton and multi-soliton solutions are obtained. Ablowitz et al. [3] implemented the inverse scattering transform method to handle the nonlinear equations of physical significance where soliton solutions and rational solutions were developed.

Some important particular cases of Eq. (1.1) are :

- Kaup-Kupershmidt equation (KK equation) [10][11][12][13][14]

$$u_t + u_{xxxxx} + 10 u u_{xxx} + 25 u_x u_{xx} + 20 u^2 u_x = 0. \tag{1.2}$$

- Sawada-Kotera equation (SK equation) [7][5]

$$u_t + u_{xxxxx} + 5uu_{xxx} + 5u_x u_{xx} + 5u^2 u_x = 0. \tag{1.3}$$

- Caudrey-Dodd-Gibbon equation

$$u_t + u_{xxxxx} + 30uu_{xxx} + 30u_x u_{xx} + 180u^2 u_x = 0. \tag{1.4}$$

- Lax equation [6]

$$u_t + u_{xxxxx} + \ 10\, uu_{xxx} + 20u_x u_{xx} + 30u^2 u_x = 0. \tag{1.5}$$

- Ito equation [8][9]

$$u_t + u_{xxxxx} + 3\, uu_{xxx} + 6u_x u_{xx} + 2\, u^2 u_x = 0. \tag{1.6}$$

As the constants α, β and γ change, the properties of the equation (1.1) drastically change. For instance, the Lax equation with $\alpha = 10$, $\beta = 20$, and $\gamma = 30$, and the SK equation where $\alpha == \beta = \gamma = 5$, are completely integrable. These two equations have N-soliton solutions and an infinite set of conserved densities. Another example is the KK equation with $\alpha = 10$, $\beta = 25$, and $\gamma = 20$, which is known to be integrable [12], and has bilinear representations [12][14], but for which the explicit form of the N-soliton solutions is not known. A fourth equation in this class is the Ito equation, with $\alpha = 3$, $\beta = 6$, and $\gamma = 2$, which is not completely integrable, but has a limited number of special conserved densities [9].

From 70's, a vast variety of simple and direct methods to find analytic solutions of nonlinear differential equations and evolution equations have been developed. Recently, the extended tanh method [1] has been successfully used in seeking the solitary wave solution and other kinds of solutions.

We will find solutions of Eq. (1.1) for $\omega \neq 0$ and $\gamma \neq 0$.

2 The extended tanh method

The extended tanh method [1] may be described in the following three steps:

Step 1. We search exact solutions of equation (1.1) in the form

$$\begin{cases} u(x,t) = v(\xi) \\ \xi = x + \lambda t, \end{cases} \tag{2.7}$$

As a result we have that the equation (1.1) is reduced to the nonlinear ordinary differential equation (ODE) of fifth order

$$\gamma v'(\xi) v(\xi)^2 + \alpha v^{(3)}(\xi) v(\xi) + \lambda v'(\xi) + \beta v'(\xi) v''(\xi) + \omega v^{(5)}(\xi) = 0 \tag{2.8}$$

Step 2. We seek solutions of (2.8) in the form

$$v(\xi) = a_0 + \sum_{i=1}^{m} \left(\varphi(\xi)^i + \varphi(\xi)^{-i} \right), \tag{2.9}$$

where

$$\varphi'(\xi) = k + \varphi^2(\xi), \tag{2.10}$$

where k is a parameter to be determined later.
The Riccati equation (2.10) has the general solutions:

- **a)** If $k < 0$: $\varphi(\xi) = -\sqrt{-k}\,\tanh(\sqrt{-k}\,\xi)$ and $\varphi(\xi) = -\sqrt{-k}\,\coth(\sqrt{-k}\,\xi)$.
- **a)** If $k = 0$: $\varphi(\xi) = -\dfrac{1}{\xi}$.
- **c)** If $k > 0$: $\varphi(\xi) = \sqrt{k}\,\tan(\sqrt{k}\,\xi)$ and $\varphi(\xi) = \sqrt{k}\,\cot(\sqrt{k}\,\xi)$.

Substituting (2.9), along with (2.10) into (2.8) and collecting all terms with the same power in $\varphi(\xi)$, we get a polynomial in the variable $\varphi = \varphi(\xi)$. This polynomial has the form

$$a\,\varphi(\xi)^{3m+1} + b\,\varphi(\xi)^{2m+3} + c\,\varphi(\xi)^{m+5} + \text{lower degree terms} \tag{2.11}$$

The parameter m can be found by balancing the high-order linear term with the nonlinear terms [2] in (2.11). We assume that $m \geq 1$ to avoid trivial solutions. The degrees of the highest terms are $m+5$ (the degree of the term $c\,\varphi(\xi)^{m+5}$), $2m+3$ (the degree of the term $b\,\varphi(\xi)^{2m+3}$) and $3m+1$ (the degree of the term $a\,\varphi(\xi)^{3m+1}$). The only integer value of m for which $3m+1 = 2m+3$ or $3m+1 = m+5$ or $2m+3 = m+5$ is $m = 2$. Equating in (2.11) the coefficients of every power of $\varphi(\xi)$ to zero, we obtain the following algebraic system in the variables k, λ, a_0, a_1, $b_1, \dots$:

- $-720b_2\omega k^5 - 24b_2^2\alpha k^3 - 12b_2^2\beta k^3 - 2b_2^3\gamma k = 0.$
- $-120b_2\omega k^5 - 30b_2b_2\alpha k^3 - 10b_2b_2\beta k^3 - 5b_2b_2^2\gamma k = 0.$
- $-1680b_2\omega k^4 - 6b_2^2\alpha k^3 - 24a_0b_2\alpha k^3 - 2b_2^2\beta k^3 - 40b_2^2\alpha k^2 - 28b_2^2\beta k^2 - 4a_0b_2^2\gamma k - 4b_2^2b_2\gamma k - 2b_2^3\gamma = 0.$
- $-240b_2\omega k^4 - 6a_0b_2\alpha k^3 - 24a_1b_2\alpha k^3 + 6a_1b_2\beta k^3 - 48b_2b_2\alpha k^2 - 22b_2b_2\beta k^2 - b_2^3\gamma k - 3a_1b_2^2\gamma k - 6a_0b_2b_2\gamma k - 5b_2b_2^2\gamma = 0.$
- $-6a_1b_2\alpha k^3 - 24a_2b_2\alpha k^3 + 2a_1b_2\beta k^3 + 8a_2b_2\beta k^3 - 1232b_2\omega k^3 - 8b_2^2\alpha k^2 - 40a_0b_2\alpha k^2 - 4b_2^2\beta k^2 - 16b_2^2\alpha k - 20b_2^2\beta k - 2a_0b_2^2\gamma k - 2a_2b_2^2\gamma k - 2a_0^2b_2\gamma k - 4a_1b_2b_2\gamma k - 2b_2\lambda k - 4a_0b_2^2\gamma - 4b_2^2b_2\gamma = 0.$
- $-\gamma b_2^3 - a_1k\gamma b_2^2 - 6a_2k^3\alpha b_2 - 8a_0k^2\alpha b_2 - 18b_2k\alpha b_2 + 2a_2k^3\beta b_2 - 14b_2k\beta b_2 - 6a_0b_2\gamma b_2 - a_0^2k\gamma b_2 - 2a_2b_2k\gamma b_2 - k\lambda b_2 - 136k^3\omega b_2 - 38a_1b_2k^2\alpha + 10a_1b_2k^2\beta - 3a_1b_2^2\gamma - 2a_0a_1b_2k\gamma = 0.$
- $-2b_2\gamma a_0^2 - 16b_2k\alpha a_0 - 2b_2^2\gamma a_0 - 6a_1b_2k^2\alpha - 24a_2b_2k^2\alpha - 2b_2^2k\alpha - 4b_2^2\beta + 2a_1b_2k^2\beta + 8a_2b_2k^2\beta - 2b_2^2k\beta - 2a_2b_2^2\gamma - 4a_1b_2b_2\gamma - 2b_2\lambda - 272b_2k^2\omega = 0.$
- $2a_1a_2\beta k^3 + 16a_1\omega k^3 + 2a_0a_1\alpha k^2 + 8a_2b_2\alpha k^2 - 2a_2b_2\beta k^2 - 16b_2\omega\, k^2 - 2a_0b_2\alpha k - 8a_1b_2\alpha k + 2a_1b_2\beta k + a_0^2a_1\gamma k + a_1^2b_2\gamma k + 2a_0a_2b_2\gamma k + 2a_1a_2b_2\gamma k + a_1\lambda k - 2b_2b_2\beta - a_1b_2^2\gamma - a_0^2b_2\gamma - 2a_0a_1b_2\gamma - 2a_2b_2b_2\gamma - b_2\lambda = 0.$

- $4a_2^2\beta k^3 + 272a_2\omega k^3 + 2a_1^2\alpha k^2 + 16a_0a_2\alpha k^2 + 2a_1^2\beta k^2 + 6a_1b_2\alpha k + 24a_2b_2\alpha k - 2a_1b_2\beta k - 8a_2b_2\beta k + 2a_0a_1^2\gamma k + 2a_0^2a_2\gamma k + 4a_1a_2b_2\gamma k + 2a_2^2b_2\gamma k + 2a_2\lambda k = 0.$
- $k\gamma a_1^3 + b_2\gamma a_1^2 + 18a_2k^2\alpha a_1 + 6b_2\alpha a_1 + 8a_0k\alpha a_1 + 14a_2k^2\beta a_1 - 2b_2\beta a_1 + a_0^2\gamma a_1 + 2a_2b_2\gamma a_1 + 6a_0a_2k\gamma a_1 + \lambda a_1 + 136k^2\omega a_1 + 38a_2b_2k\alpha - 10a_2b_2k\beta + 2a_0a_2b_2\gamma + 3a_2^2b_2k\gamma = 0.$
- $2a_2\gamma a_0^2 + 40a_2k\alpha a_0 + 2a_1^2\gamma a_0 + 4a_2^2k\gamma a_0 + 16a_2^2k^2\alpha + 6a_1b_2\alpha + 24a_2b_2\alpha + 8a_1^2k\alpha + 20a_2^2k^2\beta - 2a_1b_2\beta - 8a_2b_2\beta + 4a_1^2k\beta + 4a_1a_2b_2\gamma + 2a_2^2b_2\gamma + 4a_1^2a_2k\gamma + 2a_2\lambda + 1232a_2k^2\omega = 0.$
- $\gamma a_1^3 + 6a_0\alpha a_1 + 48a_2k\alpha a_1 + 22a_2k\beta a_1 + 6a_0a_2\gamma a_1 + 5a_2^2k\gamma a_1 + 240k\omega a_1 + 24a_2b_2\alpha - 6a_2b_2\beta + 3a_2^2b_2\gamma = 0.$
- $2k\gamma a_2^3 + 40k\alpha a_2^2 + 28k\beta a_2^2 + 4a_0\gamma a_2^2 + 24a_0\alpha a_2 + 4a_1^2\gamma a_2 + 1680k\omega a_2 + 6a_1^2\alpha + 2a_1^2\beta = 0.$
- $5a_1\gamma a_2^2 + 30a_1\alpha a_2 + 10a_1\beta a_2 + 120a_1\omega = 0.$
- $2\gamma a_2^3 + 24\alpha a_2^2 + 12\beta a_2^2 + 720\omega a_2 = 0.$

Step 3. (This is the more difficult step) Solving the previous system for k, λ, a_0, a_1, $b_1, \dots$, we get $a_1 = b_1 = 0$, so when $m = 2$ solutions have the form

$$v(\xi) = a_0 + a_2\varphi^2(\xi) + \frac{b_2}{\varphi^2(\xi)}.$$

We get the following solutions, where

$$A = 2\alpha + \beta + \sqrt{(2\alpha+\beta)^2 - 40\gamma\,\omega}, \quad B = \frac{12\gamma\,\omega - A\beta}{8\gamma} \quad \text{and} \quad C = \frac{(3A - 10\beta)\,\omega}{2A} :$$

- $a_0 = -\frac{2Ak}{\gamma}, a_2 = 0, b_0 = -\frac{3Ak^2}{\gamma}, \lambda = 16Bk^2$:

$$u_1(x,t) = -\frac{Ak}{\gamma}\left(2 + 3\cot^2\left(\sqrt{k}\ \ (x + 16Bk^2t+)\right)\right).$$

$$u_2(x,t) = \frac{Ak}{\gamma}\left(1 + 3\,\mathrm{csch}^2\left(\sqrt{-k}\,(x + 16Bk^2t)\right)\right).$$

- $a_0 = -\frac{80k\omega}{A}, a_2 = 0, b_0 = -\frac{120k^2\omega}{A}, \lambda = 16Ck^2$:

$$u_3(x,t) = -\frac{40k\omega}{A}\left(2 + 3\cot^2\left(\sqrt{k}\left(x + 256Ck^2t\right)\right)\right)$$

$$u_4(x,t) = \frac{40k\omega}{A}\left(1 + 3\,\mathrm{csch}^2\left(\sqrt{-k}\left(x + 256Ck^2t\right)\right)\right)$$

- $a_0 = -\frac{2Ak}{\gamma}, a_2 = -\frac{3A}{\gamma}, b_0 = 0, \lambda = 16Bk^2$:

$$u_5(x,t) = -\frac{Ak}{\gamma}\left(2 + 3\tan^2\left(\sqrt{k}\left(x + 16Bk^2t\right)\right)\right)$$

$$u_6(x,t) = -\frac{Ak}{\gamma}\left(2 - 3\tanh^2\left(\sqrt{-k}\left(x + 16Bk^2t\right)\right)\right).$$

- $a_0 = -\frac{80k\omega}{A}, a_2 = -\frac{120\omega}{A}, b_0 = 0, \lambda = 16Ck^2$:

$$u_7(x,t) = -\frac{40k\omega}{A}\left(2 + 3\tan^2\left(\sqrt{k}\left(x + 256Ck^2t\right)\right)\right).$$

$$u_8(x,t) = -\frac{40k\omega}{A}\left(2 - 3\tanh^2\left(\sqrt{-k}\left(x + 256Ck^2t\right)\right)\right).$$

- $a_0 = -\frac{2Ak}{\gamma}, a_2 = -\frac{3A}{\gamma}, b_2 = -\frac{3Ak^2}{\gamma}, \lambda = 256Bk^2$:

$$u_9(x,t) = -\frac{Ak}{\gamma}\left(2 + 3\tan^2\left(\sqrt{k}\left(x + 256Bk^2t\right)\right) + 3\cot^2\left(\sqrt{k}\ \left(x + 256Bk^2t\right)\right)\right).$$

$$u_{10}(x,t) = -\frac{Ak}{\gamma}\left(2 - 3\tanh^2\left(\sqrt{-k}\left(x + 256Bk^2t\right)\right) - 3\coth^2\left(\sqrt{-k}\ \left(x + 256Bk^2t\right)\right)\right).$$

- $a_0 = -\frac{80k\omega}{A}, a_2 = -\frac{120\omega}{A}, b_2 = -\frac{120k^2\omega}{A}, \lambda = 256Ck^2$:

$$u_{11}(x,t) = -\frac{40k\omega}{A}\left(2 + 3\tan^2\left(\sqrt{k}\left(x + 256Ck^2t\right)\right) + 3\cot^2\left(\sqrt{k}\left(x + 256Ck^2t\right)\right)\right).$$

$$u_{12}(x,t) = -\frac{40k\omega}{A}\left(2 - 3\tanh^2\left(\sqrt{-k}\left(x + 256Ck^2t\right)\right) - 3\coth^2\left(\sqrt{-k}\left(x + 256Ck^2t\right)\right)\right).$$

Observe that $A = 0$ only if $\omega\,\gamma = 0$.

3 Conclusions

In this paper, by using the extended tanh method and the help of a symbolic computation engine, we obtain some exact solutions for the equation (1.1). At the same time, we may find solutions for *any* particular case of this equation, for example, (1.2)-(1.6). In our opinion, this result is new in the literature.

References

[1] M.A. Abdou , A.A. Soliman, Modified extended tanh-function method and its application on nonlinear physical equations, Physics Letters A 353 (2006) 487492.

[2] E.J. Parker, B.R. Duffy, Comput. Phys. Commun. 98 (1996) 288.

[3] M.J.Ablowitz,P.A.Clarkson, Solitons, *Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University press, Cambridge,1991.

[4] R. Conte & M. Musette, *Link betwen solitary waves and projective Riccati equations*, J. Phys. A Math. **25** (1992), 5609-5623.

[5] Alvaro Salas, *Some solutions for a type of generalized Sawada-Kotera equation*, Applied Mathematics and Computation, in press, available line from july 17, 2007 at http://www.sciencedirect.com

[6] P. D. Lax, Comm. Pure Appl. Math. 21, 467-490 (1968).

[7] K. Sawada, and T. Kotera, Prog. Theor. Phys. 51, 1355-1367 (1974).

[8] M. Ito , J. Phys. Soc. Japan 49, 771-778 (1980).

[9] M. Ito, Comp. Phys. Comm. 42, 351-357 (1986).

[10] A. P. Fordy and J. Gibbons, Phys. Lett. 75 A, 325 (1980).

[11] J. Satsuma, and D.J. Kaup, J. Phys. Soc. Japan 43, 692-697 (1977).

[12] D. Kaup, Stud. Appl. Math. 62, 189-216 (1980).

[13] B. A. Kupershmidt, Phys. Lett. 102 A, 213-215 (1984).

[14] M. Jimbo, and T. Miwa, Publ. RIMS, Kyoto Univ. 19, 943-1001 (1983).